United States Patent and Trademark Office

UNITED STATES DEPARTMENT OF COMMERCE United States Patent and Trademark Office Address: COMMISSIONER FOR PATENTS P.O. Box 1450 Alexandria, Virginia 22313-1450 www.uspto.gov

APPLICATION NO.	FILING DATE	FIRST NAMED INVENTOR	ATTORNEY DOCKET NO.	CONFIRMATION NO.
10/709,355	04/29/2004	Saad Ahmed Sirohey	144482	3354
23413 7590 10/02/2007 CANTOR COLBURN, LLP 55 GRIFFIN ROAD SOUTH			EXAMINER	
			MACKOWEY, ANTHONY M	
BLOOMFIELD, CT 06002			ART UNIT	PAPER NUMBER
			2624	
			MAIL DATE	DELIVERY MODE
			10/02/2007	PAPER

Please find below and/or attached an Office communication concerning this application or proceeding.

The time period for reply, if any, is set in the attached communication.

	A A.				
•	Application No.	Applicant(s)			
Óssica Action O	10/709,355	SIROHEY ET AL.			
Office Action Summar	Examiner	Art Unit			
•	Anthony Mackowey	2624			
eriod for Reply .	munication appears on the cover sheet wi				
 WHICHEVER IS LONGER, FROM The Extensions of time may be available under the provafter SIX (6) MONTHS from the mailing date of this If NO period for reply is specified above, the maximer Failure to reply within the set or extended period for the set or extended period fo	num statutory period will apply and will expire SIX (6) MON reply will, by statute, cause the application to become AB on the mailing date of this communication, even if the state of the communication.	CATION. Teply be timely filed ITHS from the mailing date of this communication. BANDONED (35 U.S.C. & 133)			
atus		•			
1) Responsive to communication(s	s) filed on 02 July 2007.				
2a) This action is FINAL.					
3) Since this application is in cond	ition for allowance except for formal matte	ers, prosecution as to the merits is			
	ractice under <i>Ex parte Quayle</i> , 1935 C.D				
sposition of Claims	•	•			
4)⊠ Claim(s) <u>1-7 and 9-19</u> is/are per	nding in the application.				
•	is/are withdrawn from consideration.				
5) Claim(s) is/are allowed.		·			
6)⊠ Claim(s) <u>1-7 and 9-19</u> is/are reje	ected.				
7) Claim(s) is/are objected to	(O.				
8) Claim(s) are subject to re	estriction and/or election requirement.				
plication Papers					
9) The specification is objected to b	ov the Examiner.				
-	2004 is/are: a)⊠ accepted or b)□ objec	ted to by the Examiner.			
	objection to the drawing(s) be held in abeyan	-			
	uding the correction is required if the drawing(
e	ed to by the Examiner. Note the attached	• •			
iority under 35 U.S.C. § 119		•			
12) Acknowledgment is made of a cl	aim for foreign priority under 35 U.S.C. § of:	119(a)-(d) or (f).			
· · · · · · · · · · · · · · · · · · ·	ority documents have been received.	•			
2. Certified copies of the price	prity documents have been received in Ap	pplication No			
3. Copies of the certified cop	pies of the priority documents have been	received in this National Stage			
•	national Bureau (PCT Rule 17.2(a)).	•			
* See the attached detailed Office a	action for a list of the certified copies not	received.			
achment(s)		•			
Notice of References Cited (PTO-892)		Summary (PTO-413)			
Notice of Draftsperson's Patent Drawing Review		s)/Mail Date Informal Patent Application			
Information Disclosure Statement(s) (PTO/SB Paper No(s)/Mail Date	(708) 5) Notice of in	• •			

Art Unit: 2624

DETAILED ACTION

Response to Arguments

Applicant's arguments, see page 1, filed July 2, 2007, with respect to the objections to the claims have been fully considered and are persuasive. The objections to the claims have been withdrawn.

Applicant's arguments, see page 2, filed July 2, 2007, with respect to the objections to the specification have been fully considered and are persuasive. The objections to the specification have been withdrawn.

Applicant's arguments, see page 2, filed July 2, 2007, with respect to the rejection of claims 4 and 7 under 35 U.S.C. 112, second paragraph have been fully considered and are persuasive. The rejection of claims 4 and 7 under 35 U.S.C. 112, second paragraph has been withdrawn.

Applicant's arguments filed July 2, 2007, with respect to the claim rejections under 35 U.S.C. 103(a) have been fully considered but they are not persuasive. Applicant submits, "Limerely discloses classifying nodule candidates in a region of interest and filtering an entire set of image data. Therefore, Li cannot disclose or suggest 'isolating a selected region of interest from the multidimensional dataset...generating a plurality of differential operators for the selected region of interest using discrete approximation of an analytic function' as recited in independent claim 1." Examiner agrees with applicant in what Li explicitly recites as a "region of interest" (col. 12, lines 35-55) does not appear to be filtered with the line and plane enhancement filters (col. 12, lines 56-67). However, Li teaches the lung regions are first segmented from the original

Art Unit: 2624

images and then the enhancement filters are applied to the segmented lung regions (Fig. 2; col. 12, lines 13-34). While the explicitly recited "region of interest" of Li does not meet the limitations recited in claim 1, the segmentation of the lung region and subsequent filtering meet the requirements of the region of interest recited in claim 1 (and similarly in claims 15, 16, 17, 18 and 19). Therefore, the segmentation of the lung regions taught by Li may be reasonably construed as isolating a selected region of interest as recited in the claims. Examiner notes that claim 10 recites, "wherein said isolating a selected region of interest further includes isolating lung tissue for a pair of lungs" and the Examiner had indicated in the rejection of claim 10 in the previous Office action Li taught such a limitation (see page 17 of Office action mailed May 1, 2007). Therefore, because Li is not deficient in the teaching of the isolation of a selected region of interest from which a plurality of differential operators are generated, the combinations of the teachings of Li, Lindeberg and Hu disclose or suggest all the limitations of the claims.

Claim Rejections - 35 USC § 103

The following is a quotation of 35 U.S.C. 103(a) which forms the basis for all obviousness rejections set forth in this Office action:

(a) A patent may not be obtained though the invention is not identically disclosed or described as set forth in section 102 of this title, if the differences between the subject matter sought to be patented and the prior art are such that the subject matter as a whole would have been obvious at the time the invention was made to a person having ordinary skill in the art to which said subject matter pertains. Patentability shall not be negatived by the manner in which the invention was made.

Claims 1-7, 11, 12 and 14-19 are rejected under 35 U.S.C. 103(a) as being unpatentable over US 6,937,776 to Li et al. (hereafter referred to as "Li") in view of "Discrete Derivative Approximations with Scale-Space Properties: A Basis for Low-Level Feature Extraction" by Lindeberg.

· Art Unit: 2624

Regarding claim 1, Li discloses a method for processing of a multi-dimensional dataset corresponding to an imaging volume (col. 4, lines 59-67), the method comprising:

accessing the multi-dimensional dataset (col. 19, lines 29-40);

isolating a selected region of interest from the multidimensional dataset, said selected region of interest comprising a subset of the imaging volume (col. 12, lines 12-34);

generating a plurality of differential operators for selected region of interest using an approximation of an analytic function (col. 7, line 11 – col. 10, line 37; col. 12, lines 30-67, Li teaches generating the second order derivatives of the images and approximations to the second order derivatives and constructing a Hessian matrix); and

forming a plurality of geometric responses based on a plurality of differential operators resultant from said generating (col. 6, line 10 – col. 7, line 32; col. 12, lines 30-67, Li teaches geometric filters based on the second order derivatives.).

Li discloses generating a plurality of differential operators for the multi-dimensional dataset using an approximation of an analytic function (col. 10, line 6-37) but does not explicitly disclose using a discrete approximation of an analytic function. However, Lindeberg clearly teaches using the discrete approximations of an analytic function (discrete derivative approximations) to determine the second order derivatives for use in a feature detector (Abstract; page 19, Section 6.1.3, second paragraph; page 16, Figure 3).

The teachings of Li and Lindeberg are combinable because they are both concerned with using filters derived from second order derivatives for feature extraction. It would have been obvious to one of ordinary skill in the art that the method taught by Li could be modified so the generating of a plurality of differential operators for the multi-dimensional dataset uses a discrete

Art Unit: 2624

approximation of an analytic function as taught by Lindeberg because the discrete approximations can be computed directly from smoothed image values at different scales without the need for repeating the smoothing operation, thus reducing the computation requirements making it more efficient (Lindeberg, page 2, lines 32-41 (last full paragraph); page 12, section 4.1).

Regarding claim 2, Lindeberg further discloses scale-space processing the multidimensional dataset with multi-resolution sampling (page 25, section 7.1).

Regarding claim 3, Li further discloses iterating said generating and forming over several scales to determine said plurality of responses for each scale; and determining said plurality of geometric responses based on said iterating (Fig. 11; col. 11, lines 29-60).

Regarding claim 4, Li further discloses filtering the multi-dimensional dataset with a smoothing kernel based on an analytic function; said smoothing kernel generating a filtered multi-dimensional dataset (Fig. 11, S3; col. 10, line 46 – col. 11, line 60).

Regarding claim 5, Li further discloses said analytic function is a Gaussian (Fig. 11, S3; col. 10, line 46 – col. 11, line 60).

Art Unit: 2624

Regarding claim 6, Li further discloses said plurality of differential operators correspond to an n-th derivative of said analytic function, where n is greater than or equal to one (col. 7, line 11 – col. 11, line 60, Li teaches second derivatives.)

Regarding claim 7, the combination of Li and Lindeberg further discloses identifying a plurality of discrete derivative approximations that when convolved with said analytic function, approximates an analytical derivative of said analytic function. Lindeberg teaches identifying a plurality discrete approximations and Li teaches identifying approximations that when convolved with said analytic function, approximate an analytic derivative of said analytic function (see arguments and citations presented above for claim 1 and additionally Li, col. 10, line 38 - col. 11, line 7).

Li and Lindeberg are silent with regard to optimizing said discrete derivative approximations in a least squares sense to reduce an error between said plurality of discrete derivative approximations and said analytical derivative of said analytic function. The Examiner takes Official Notice that optimization in a least squares sense if well known in the art and it would have been obvious to one of ordinary skill in the art to optimize the discrete derivative approximations in a least squares sense to reduce an error between the discrete derivative approximations and the analytical derivative of the analytic function because optimization using least squares techniques is exceedingly well known for quantifying and reducing differences (error) between two functions.

Art Unit: 2624

Regarding claim 11, Lindeberg further discloses generating a downsampled multidimensional dataset based on said multi-resolution sampling (page 25, Section 7.1).

Regarding claim 12, Li further discloses isolating a selected region of interest from at least one of said multi-dimensional dataset and said downsampled multi-dimensional dataset, said selected region of interest comprising a subset of the imaging volume (col. 12, lines 12-55, Examiner notes the use of alternative language. Li teaches isolating a selected region of interest from the multi-dimensional dataset.).

Regarding claim 14, neither Li nor Lindeberg explicitly disclose said processing of a multi-dimensional dataset is executed in less than one minute, however one of ordinary skill in the art at the time the invention was made would have found it obvious that the processing taught by the combination of Li and Lindeberg, as presented above in the rejection of claim 1, is capable of being performed in less than one minute. If the presently claimed processing method is capable of being performed in less than one minute, because the combination of Li and Lindeberg disclose the same steps, then the processing method taught by the combination of Li and Lindeberg would also be performed in less than one minute when performed by an equivalent processing apparatus/computer.

Regarding claim 15, Li discloses a method for processing of a multi-dimensional dataset corresponding to an imaging volume (col. 4, lines 59-67).

Art Unit: 2624

Li does not disclose processing the multidimensional dataset with multi-resolution sampling to establish a downsampled multidimensional dataset, however Lindeberg discloses multi-resolution sampling to establish a downsampled multidimensional dataset (page 25, section 7.1). One of ordinary skill in the art at the time the invention was made would have been motivated to modify the method taught by Li to include multiresolution sampling to establish a downsampled multidimensional dataset as taught by Lindeberg in order to improve computational efficiency (Lindeberg, page 25, section 7.1, first paragraph).

Li further discloses identifying a region of interest from the multi-dimensional dataset; said region of interest comprising a subset of the imaging volume (col. 12, lines 12-34).

In view of the modification to the method of Li to include sampling the multi-resolution sampling as taught by Lindeberg is would have further been obvious to one of ordinary skill in the art to process said downsampled multidimensional dataset based on said region of interest and establishing a multi-dimensional datasubset in view of Li's teaching of establishing a datasubset based on a region of interest (col. 12, lines 12-34).

Li further discloses filtering the a multi-dimensional datasubset with a smoothing kernel based on an analytic function; said smoothing kernel generating a filtered multi-dimensional datasubset (Fig. 11, S3; col. 10, line 46 – col. 11, line 60).

Li also discloses generating a plurality of differential operators for the multi-dimensional datasubset using an approximation of an analytic function (col. 10, line 6-37) but does not explicitly disclose using a discrete approximation of an analytic function. However, Lindeberg clearly teaches using the discrete approximations of an analytic function (discrete derivative approximations) to determine the second order derivatives for use in a feature detector (Abstract;

Art Unit: 2624

page 19, Section 6.1.3, second paragraph; page 16, Figure 3). It would have been obvious to one of ordinary skill in the art that the method taught by Li could be modified so the generating of a plurality of differential operators for the multi-dimensional dataset uses a discrete approximation of an analytic function as taught by Lindeberg because the discrete approximations can be computed directly from smoothed image values at different scales without the need for repeating the smoothing operation, thus reducing the computation requirements making it more efficient (Lindeberg, page 2, lines 32-41 (last full paragraph); page 12, section 4.1).

Li further discloses forming a plurality of geometric responses based on a plurality of differential operators resultant from said generating (col. 6, line 10 – col. 7, line 32).

Regarding claim 16, Li discloses a method for processing of a multi-dimensional dataset (col. 4, lines 59-67). Li does not disclose processing the multi-dimensional dataset in a multi-resolution framework. However, Lindeberg discloses multi-resolution sampling of a multidimensional dataset (page 25, section 7.1). One of ordinary skill in the art at the time the invention was made would have been motivated to modify the method taught by Li to include multiresolution process the multi-dimensional dataset in a multi-resolution framework as taught by Lindeberg in order to improve computational efficiency (Lindeberg, page 25, section 7.1, first paragraph).

Li further discloses isolating a selected region of interest from said multidimensional dataset and establishing a multidimensional datasubset, said selected region of interest comprising a subset of the imaging volume (col. 12, lines 12-34);

Art Unit: 2624

convolving said multidimensional datasubset with an analytic function to obtain a first convolution product (Fig. 11; col. 10, line 46 – col. 11, line 60); and

determining a plurality of derivative approximations to an analytic function (col. 10, line 6-37).

Li does not explicitly disclose determining a plurality of discrete derivative approximations to an analytic function. However, Lindeberg clearly teaches using the discrete approximations of an analytic function (discrete derivative approximations) to determine the second order derivatives for use in a feature detector (Abstract; page 19, Section 6.1.3, second paragraph; page 16, Figure 3). It would have been obvious to one of ordinary skill in the art that the method taught by Li could be modified to include determining a plurality of discrete derivative approximations to an analytic function as taught by Lindeberg because the discrete approximations can be computed directly from smoothed image values at different scales without the need for repeating the smoothing operation, thus reducing the computation requirements making it more efficient (Lindeberg, page 2, lines 32-41 (last full paragraph); page 12, section 4.1).

Li and Lindeberg are silent with regard to optimizing said discrete derivative approximations in a least squares sense to reduce an error between said plurality of discrete derivative approximations and said analytical derivative of said analytic function. The Examiner takes Official Notice that optimization in a least squares sense if well known in the art and it would have been obvious to one of ordinary skill in the art to optimize the discrete derivative approximations in a least squares sense to reduce an error between the discrete derivative approximations and the analytical derivative of the analytic function because optimization using

Art Unit: 2624

least squares techniques is exceedingly well known for quantifying and reducing differences (error) between two functions.

The combination of Li and Lindeberg further discloses convolving said first convolution product with the plurality of discrete approximations of partial derivatives of an analytic function to create a plurality of second convolution products (Li, Fig. 11; col. 10, line 46 – col. 11, line 60);

forming a plurality of Hessian matrices from said plurality of second convolution products (Li, Fig. 11; col. 11, line 44 – col. 12, line 11);

determining a plurality of eigenvalue decompositions for said plurality of said Hessian matrices (col. 9, line 63 – col. 10, line 5); and

combining eigenvalues resultant from said decompositions to represent spherical and cylindrical responses to elements of said multidimensional datasubset (Li, col. 6, line 10 – col. 9, line 62).

Regarding claim 17, Li further discloses a system for processing of a multi-dimensional dataset corresponding to an imaging volume (col. 4, lines 59-67; col. 18, lines 37-44), the system comprising:

a means for accessing the multi-dimensional dataset (col. 18, lines 37-44; col. 19, lines 29-40);

a means for isolating a selected region of interest from the multi-dimensional dataset, said selected region of interest comprising a subset of the imaging volume(col. 18, lines 37-44; col. 12, lines 12-34)

Art Unit: 2624

a means for generating a plurality of differential operators for selected region of interest using an approximation of an analytic function (col. 18, lines 37-44; col. 7, line 11 - col. 10, line 37; col. 12, lines 30-67); and

a means for forming a plurality of geometric responses based on a plurality of differential operators resultant from said generating (col. 18, lines 37-44; col. 6, line 10 – col. 7, line 32; col. 12, lines 30-67).

Li discloses generating a plurality of differential operators for the multi-dimensional dataset using approximations of an analytic function (col. 10, line 6-37) but does not explicitly disclose using a discrete approximation of an analytic function. However, Lindeberg clearly teaches using the discrete approximations of an analytic function (discrete derivative approximations) to determine the second order derivatives for use in a feature detector (Abstract; page 19, Section 6.1.3, second paragraph; page 16, Figure 3).

The teachings of Li and Lindeberg are combinable because they are both concerned with using filters derived from second order derivatives for feature extraction. It would have been obvious to one of ordinary skill in the art that the system taught by Li could be modified so the generating of a plurality of differential operators for the multi-dimensional dataset uses a discrete approximation of an analytic function as taught by Lindeberg because the discrete approximations can be computed directly from smoothed image values at different scales without the need for repeating the smoothing operation, thus reducing the computation requirements making it more efficient (Lindeberg, page 2, lines 32-41 (last full paragraph); page 12, section 4.1).

Art Unit: 2624

Regarding claim 18, Li discloses a system for processing of a multi-dimensional dataset corresponding to an imaging volume (col. 4, lines 59-67), the system comprising:

an imaging system comprising;

a radiation source configured to generate a radiation beam incident upon an object,
a radiation detector, said radiation detector configured to receive an attenuated radiation
beam responsive to said radiation beam incident upon said object and produce an electrical signal
responsive to an intensity of attenuated radiation beam, and

wherein said radiation source and said radiation detector disposed about an object cavity.

Li discloses the multi-dimensional dataset can be obtained from an X-ray CT apparatus (col. 19, lines 29-31; col. 12, lines 12-34). The above limitations are inherent to a conventional X-ray CT apparatus being used to image an object.

Li further discloses a processing device in operable communication with said radiation detector configured to execute a method for processing of a multi-dimensional dataset corresponding to an imaging volume (col. 18, lines 37-44; col. 19, lines 29-40), the method comprising;

accessing the multi-dimensional dataset (col. 19, lines 29-40),

isolating a selected region of interest from the multi-dimensional dataset, said selected region of interest comprising a subset of the imaging volume (col. 12, lines 12-34),

generating a plurality of differential operators for selected region of interest using an approximation of an analytic function (col. 7, line 11 – col. 10, line 37; col. 12, lines 30-67), and

forming a plurality of geometric responses based on a plurality of differential operators resultant from said generating (col. 6, line 10 - col. 7, line 32; col. 12, lines 30-67).

Art Unit: 2624

Li discloses generating a plurality of differential operators for the multi-dimensional dataset using an approximation of an analytic function (col. 10, line 6-37) but does not explicitly disclose using a discrete approximation of an analytic function. However, Lindeberg clearly teaches using the discrete approximations of an analytic function (discrete derivative approximations) to determine the second order derivatives for use in a feature detector (Abstract; page 19, Section 6.1.3, second paragraph; page 16, Figure 3).

The teachings of Li and Lindeberg are combinable because they are both concerned with using filters derived from second order derivatives for feature extraction. It would have been obvious to one of ordinary skill in the art that the method taught by Li could be modified so the generating of a plurality of differential operators for the multi-dimensional dataset uses a discrete approximation of an analytic function as taught by Lindeberg because the discrete approximations can be computed directly from smoothed image values at different scales without the need for repeating the smoothing operation, thus reducing the computation requirements making it more efficient (Lindeberg, page 2, lines 32-41 (last full paragraph); page 12, section 4.1).

Regarding claim 19, Li discloses a computer data storage device, said computer data storage device including computer readable program code, the computer readable program code comprising a method for processing of a multi-dimensional dataset corresponding to an imaging volume (col. 18, line 45 – col. 19, line 20). Regarding the method, arguments analogous to those presented above for claim 1 are applicable to claim 19.

Art Unit: 2624

Claims 9 and 10 are rejected under 35 U.S.C. 103(a) as being unpatentable over the combination of Li and Lindeberg as applied to claim 1 above, and further in view of "Automatic Lung Segmentation for Accurate Quantitation of Volumetric X-Ray CT Images" by Hu et al. (hereafter "Hu").

Regarding claim 9, Li further discloses said isolating a selected region of interest includes image threshold filtering configured to eliminate selected portions of the imaging volume (col. 12, lines 12-55) but is silent with regard to a morphology process. However, Hu teaches a lung segmentation method including morphological processes (page 493-494, Section C).

The teachings of Li and Hu are combinable because they are both concerned with segmenting lung regions from CT images. It would have been obvious to one of ordinary skill in the art at the time the invention was made for the isolating of the region of interest taught by Li to include morphological processing as taught by Hu so the automatic isolation/segmentation more closely mimics segmentation performed manually (page 493, Section C, second paragraph).

Regarding claim 10, Li further discloses said isolating a selected region of interest further includes isolating lung tissue for a pair of lungs comprising filtering with a threshold algorithm (col. 12, lines 12-55) but does not explicitly disclose the steps recited in claim 10.

Hu discloses:

filtering with a high threshold algorithm to isolate solid tissue and bone (page 491, Section A.1);

Art Unit: 2624

filling holes with a three-dimensional hole-filling algorithm to fill a portion of remain contained inside said solids (page 491, Section A.2);

filtering with a low threshold algorithm to isolate parenchyma of a pair of lungs from the solid tissue and bone (page 491, Section A.1);

splitting and isolating said pair of lungs with a morphology erosion algorithm (page 491-493, Section B;

closing and filing airways and vascular structures entering said pair of lungs with a morphology closure algorithm (page 493, Section C.1); and

filling remaining holes with a three-dimensional hole-filling algorithm to yield another multidimensional dataset corresponding to the selected region of interest (page 494, Sections 4 and 5; Fig. 3).

The teachings of Li and Hu are combinable because they are both concerned with segmenting lung regions from CT images. It would have been obvious to one of ordinary skill in the art at the time the invention was made for the isolating of the region of interest taught by Li to include steps recited claim 10 as taught by Hu so the automatic isolation/segmentation more closely mimics segmentation performed manually (page 493, Section C, second paragraph).

Claim 13 is rejected under 35 U.S.C. 103(a) as being unpatentable over the combination of Li and Lindeberg as applied to claim 11 above, and further in view of "Automatic Lung Segmentation for Accurate Quantitation of Volumetric X-Ray CT Images" by Hu et al. (hereafter "Hu").

Art Unit: 2624

Regarding claim 13, Li further discloses said isolating a selected region of interest includes image threshold filtering configured to eliminate selected portions of the imaging volume (col. 12, lines 12-34) but is silent with regard to a morphology process. However, Hu teaches a lung segmentation method including morphological processes (page 493-494, Section C).

The teachings of Li and Hu are combinable because they are both concerned with segmenting lung regions from CT images. It would have been obvious to one of ordinary skill in the art at the time the invention was made for the isolating of the region of interest taught by Li to include morphological processing as taught by Hu so the automatic isolation/segmentation more closely mimics segmentation performed manually (page 493, Section C, second paragraph).

Conclusion

Applicant's amendment necessitated the new ground(s) of rejection presented in this Office action. Accordingly, **THIS ACTION IS MADE FINAL**. See MPEP § 706.07(a). Applicant is reminded of the extension of time policy as set forth in 37 CFR 1.136(a).

A shortened statutory period for reply to this final action is set to expire THREE MONTHS from the mailing date of this action. In the event a first reply is filed within TWO MONTHS of the mailing date of this final action and the advisory action is not mailed until after the end of the THREE-MONTH shortened statutory period, then the shortened statutory period will expire on the date the advisory action is mailed, and any extension fee pursuant to 37 CFR 1.136(a) will be calculated from the mailing date of the advisory action. In no event,

Art Unit: 2624

however, will the statutory period for reply expire later than SIX MONTHS from the date of this final action.

Applicant's amendment necessitated the new ground(s) of rejection presented in this Office action. Accordingly, **THIS ACTION IS MADE FINAL**. See MPEP § 706.07(a). Applicant is reminded of the extension of time policy as set forth in 37 CFR 1.136(a).

A shortened statutory period for reply to this final action is set to expire THREE MONTHS from the mailing date of this action. In the event a first reply is filed within TWO MONTHS of the mailing date of this final action and the advisory action is not mailed until after the end of the THREE-MONTH shortened statutory period, then the shortened statutory period will expire on the date the advisory action is mailed, and any extension fee pursuant to 37 CFR 1.136(a) will be calculated from the mailing date of the advisory action. In no event, however, will the statutory period for reply expire later than SIX MONTHS from the date of this final action.

Any inquiry concerning this communication or earlier communications from the examiner should be directed to Anthony Mackowey whose telephone number is (571) 272-7425. The examiner can normally be reached on M-F 9:00-6:00.

If attempts to reach the examiner by telephone are unsuccessful, the examiner's supervisor, Bella Matthew can be reached on (571) 272-7778. The fax phone number for the organization where this application or proceeding is assigned is 571-273-8300.

Art Unit: 2624

Information regarding the status of an application may be obtained from the Patent Application Information Retrieval (PAIR) system. Status information for published applications may be obtained from either Private PAIR or Public PAIR. Status information for unpublished applications is available through Private PAIR only. For more information about the PAIR system, see http://pair-direct.uspto.gov. Should you have questions on access to the Private PAIR system, contact the Electronic Business Center (EBC) at 866-217-9197 (toll-free). If you would like assistance from a USPTO Customer Service Representative or access to the automated information system, call 800-786-9199 (IN USA OR CANADA) or 571-272-1000.

AM 9/27/07

ANDREW W. JOHNS